

Algorithms and Data Structures

The Simplex Method

Linear Programs in Standard Form

$$\begin{array}{ll}\text{maximise} & \sum_{j=1}^n c_j x_j \\ \text{subject to} & \sum_{j=1}^n \alpha_{ij} x_j \leq b_i, \quad i = 1, \dots, m \\ & x_j \geq 0, \quad j = 1, \dots, n\end{array}$$

Linear Programs in Standard Form

maximise $\sum_{j=1}^n c_j x_j$

subject to $\sum_{j=1}^n \alpha_{ij} x_j \leq b_i, \quad i = 1, \dots, m$

$$x_j \geq 0, \quad j = 1, \dots, n$$

Linear Programs in Standard Form

$$\begin{array}{ll}\text{maximise} & \sum_{j=1}^n c_j x_j \\ \text{subject to} & \sum_{j=1}^n \alpha_{ij} x_j \leq b_i, \quad i = 1, \dots, m \\ & x_j \geq 0, \quad j = 1, \dots, n\end{array}$$

The Linear Programming problem

The Linear Programming problem

Given a Linear Program (LP) in standard form:

The Linear Programming problem

Given a Linear Program (LP) in standard form:

Return an **optimal solution** (i.e., a feasible solution that maximises the objective function), or

The Linear Programming problem

Given a Linear Program (LP) in standard form:

Return an **optimal solution** (i.e., a feasible solution that maximises the objective function), or

Return that the LP is **infeasible** , or

The Linear Programming problem

Given a Linear Program (LP) in standard form:

Return an **optimal solution** (i.e., a feasible solution that maximises the objective function), or

Return that the LP is **infeasible** , or

Return that the LP is **unbounded**.

The Simplex Method (explained via example)

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$\begin{aligned} 2x_1 + 3x_2 + x_3 &\leq 5 \\ 4x_1 + x_2 + 2x_3 &\leq 11 \\ 3x_1 + 4x_2 + 2x_3 &\leq 8 \\ x_1, x_2, x_3 &\geq 0 \end{aligned}$$

Step 1: Slack Variables

Step 1: Slack Variables

For each constraint we introduce a *slack variable*:

Step 1: Slack Variables

For each constraint we introduce a *slack variable*:

e.g., for the constraint $2x_1 + 3x_2 + x_3 \leq 5$, we introduce variable w_1 and we write

$$w_1 = 5 - 2x_1 - 3x_2 - x_3$$

Step 1: Slack Variables

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$2x_1 + 3x_2 + x_3 \leq 5$$
$$4x_1 + x_2 + 2x_3 + 3 \leq 11$$
$$3x_1 + 4x_2 + 2x_3 \leq 8$$
$$x_1, x_2, x_3 \geq 0$$

Step 1: Slack Variables

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$w_1 = 5 - 2x_1 + 3x_2 + x_3$$
$$w_2 = 11 - 4x_1 + x_2 + 2x_3 + 3$$
$$w_3 = 8 - 3x_1 + 4x_2 + 2x_3$$
$$x_1, x_2, x_3 \geq 0$$

Step 1: Slack Variables

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$\begin{aligned}w_1 &= 5 - 2x_1 + 3x_2 + x_3 \\w_2 &= 11 - 4x_1 + x_2 + 2x_3 + 3 \\w_3 &= 8 - 3x_1 + 4x_2 + 2x_3 \\x_1, x_2, x_3 &\geq 0\end{aligned}$$

Is this equivalent to the original LP?

Step 1: Slack Variables

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$w_1 = 5 - 2x_1 + 3x_2 + x_3$$
$$w_2 = 11 - 4x_1 + x_2 + 2x_3 + 3$$
$$w_3 = 8 - 3x_1 + 4x_2 + 2x_3$$
$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Step 1: Slack Variables

For each constraint we introduce a *slack variable*:

e.g., for the constraint $2x_1 + 3x_2 + x_3 \leq 5$, we introduce variable w_1 and we write

$$w_1 = 5 - 2x_1 - 3x_2 - x_3$$

Step 1: Slack Variables

For each constraint we introduce a *slack variable*:

e.g., for the constraint $2x_1 + 3x_2 + x_3 \leq 5$, we introduce variable w_1 and we write

$$w_1 = 5 - 2x_1 - 3x_2 - x_3$$

We also introduce a slack variable ζ for the objective function.

Step 1: Slack Variables

Maximise $\zeta = 5x_1 + 4x_2 + 3x_3$

subject to

$$w_1 = 5 - 2x_1 + 3x_2 + x_3$$
$$w_2 = 11 - 4x_1 + x_2 + 2x_3 + 3$$
$$w_3 = 8 - 3x_1 + 4x_2 + 2x_3$$
$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Dictionaries

Maximise $\zeta =$ $+5 \ x_1$ $+4 \ x_2$ $+3 \ x_3$

subject to

$w_1 = 5$	$-2 \ x_1$	$-3 \ x_2$	$- \ x_3$
$w_2 = 11$	$-4 \ x_1$	$- \ x_2$	$-2 \ x_3$
$w_3 = 8$	$-3 \ x_1$	$-4 \ x_2$	$-2 \ x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables



nonbasic variables

Step 1: Slack Variables

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$w_1 = 5 - 2x_1 + 3x_2 + x_3$$
$$w_2 = 11 - 4x_1 + x_2 + 2x_3 + 3$$
$$w_3 = 8 - 3x_1 + 4x_2 + 2x_3$$
$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

The Simplex Method (strategy)

The Simplex Method (strategy)

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

The Simplex Method (strategy)

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

Improve this solution to some $\bar{x}_1, \bar{x}_2, \bar{x}_3, \bar{w}_1, \bar{w}_2, \bar{w}_3$ such that
 $5\bar{x}_1 + 4\bar{x}_2 + 3\bar{x}_3 > 5x_1 + 4x_2 + 3x_3$

The Simplex Method (strategy)

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

Improve this solution to some $\bar{x}_1, \bar{x}_2, \bar{x}_3, \bar{w}_1, \bar{w}_2, \bar{w}_3$ such that
 $5\bar{x}_1 + 4\bar{x}_2 + 3\bar{x}_3 > 5x_1 + 4x_2 + 3x_3$

Continue until no further improvement is possible (in that case we are at an optimal solution).

The Simplex Method (strategy)

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

Improve this solution to some $\bar{x}_1, \bar{x}_2, \bar{x}_3, \bar{w}_1, \bar{w}_2, \bar{w}_3$ such that
 $5\bar{x}_1 + 4\bar{x}_2 + 3\bar{x}_3 > 5x_1 + 4x_2 + 3x_3$

Continue until no further improvement is possible (in that case we are at an optimal solution).

Does this remind you of something?

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$



nonbasic variables

basic variables

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$



nonbasic variables

basic variables

Start with a feasible solution $x_1, x_2, x_3, w_1, w_2, w_3$

Suggestions?

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables



nonbasic variables

$x_1 = x_2 = x_3 = 0$

Dictionaries

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$-2 x_1 \quad -3 x_2 \quad - x_3$

$-4 x_1 \quad - x_2 \quad -2 x_3$

$-3 x_1 \quad -4 x_2 \quad -2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$



nonbasic variables

basic variables

$x_1 = x_2 = x_3 = 0$

$w_1 = 5, w_2 = 11, w_3 = 8$


Dictionaries


Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

 **basic variables**

 **nonbasic variables**

A solution obtained by setting all the nonbasic variables to 0 is called a **basic feasible solution**.

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0 \qquad w_1 = 5, w_2 = 11, w_3 = 8$$

We can increase the value of some nonbasic variable, e.g., x_1

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0 \qquad w_1 = 5, w_2 = 11, w_3 = 8$$

We can increase the value of some nonbasic variable, e.g., x_1

We should not violate any constraints though!

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0 \qquad w_1 = 5, w_2 = 11, w_3 = 8$$

We can increase the value of some nonbasic variable, e.g., x_1

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 \ x_1$ $+4 \ x_2$ $+3 \ x_3$

subject to

$w_1 = 5$	$-2 \ x_1$	$-3 \ x_2$	$- \ x_3$
$w_2 = 11$	$-4 \ x_1$	$- \ x_2$	$-2 \ x_3$
$w_3 = 8$	$-3 \ x_1$	$-4 \ x_2$	$-2 \ x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For w_1 , x_1 can become as large as $5/2 = 30/12$.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For w_1 , x_1 can become as large as $5/2 = 30/12$.

For w_2 , x_1 can become as large as $11/4 = 33/12$.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For w_1 , x_1 can become as large as $5/2 = 30/12$.

For w_2 , x_1 can become as large as $11/4 = 33/12$.

For w_3 , x_1 can become as large as $32/12$.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 5/2, \quad x_2 = x_3 = 0$$

For w_1 , x_1 can become as large as $5/2 = 30/12$.

For w_2 , x_1 can become as large as $11/4 = 33/12$.

For w_3 , x_1 can become as large as $32/12$.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

For w_1 , x_1 can become as large as $5/2 = 30/12$.

For w_2 , x_1 can become as large as $11/4 = 33/12$.

For w_3 , x_1 can become as large as $32/12$.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \qquad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \qquad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \qquad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ $-2 x_1$ $-3 x_2$ $- x_3$
 $w_2 = 11$ $-4 x_1$ $- x_2$ $-2 x_3$
 $w_3 = 8$ $-3 x_1$ $-4 x_2$ $-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ leaving variable

$w_2 = 11$

$w_3 = 8$

$-2 x_1$	$-3 x_2$	$- x_3$
$-4 x_1$	$- x_2$	$-2 x_3$
$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ $-2 x_1 - 3 x_2 - x_3$ } just rearranging
 $w_2 = 11$ $-4 x_1 - x_2 - 2 x_3$
 $w_3 = 8$ $-3 x_1 - 4 x_2 - 2 x_3$

leaving variable

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ leaving variable

$w_2 = 11$

$w_3 = 8$

$-2 x_1$	$-3 x_2$	$- x_3$
$-4 x_1$	$- x_2$	$-2 x_3$
$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

} just rearranging

} what about here?

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ leaving variable

$w_2 = 11$

$w_3 = 8$

$-2 x_1$	$-3 x_2$	$- x_3$
$-4 x_1$	$- x_2$	$-2 x_3$
$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

} just rearranging

} what about here?

} "row operations"

$$x_1 = 5/2, \quad x_2 = x_3 = 0 \quad w_1 = 0, \quad w_2 = 1, \quad w_3 = 1/2$$

We need to rearrange the inequalities, so that x_1 now only appears on the left.

This gives rise to a new dictionary, where x_1 is now **basic** and w_1 is **nonbasic**.

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to $w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$$-2 x_1 \quad -3 x_2 \quad - x_3$$

$$-4 x_1 \quad - x_2 \quad -2 x_3$$

$$-3 x_1 \quad -4 x_2 \quad -2 x_3$$

} just rearranging
} what about here?
} "row operations"

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to $w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$$-2 x_1 \quad -3 x_2 \quad - x_3$$

$$-4 x_1 \quad - x_2 \quad -2 x_3$$

$$-3 x_1 \quad -4 x_2 \quad -2 x_3$$

} just rearranging

} what about here?

} “row operations”

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Notice that $w_2 - 2w_1 = 11 - 4x_1 - x_2 - 2x_3 - 10 + 4x_1 + 6x_2 + 2x_3$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to $w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$$-2 x_1 \quad -3 x_2 \quad - x_3$$

$$-4 x_1 \quad - x_2 \quad -2 x_3$$

$$-3 x_1 \quad -4 x_2 \quad -2 x_3$$

} just rearranging

} what about here?

} “row operations”

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Notice that $w_2 - 2w_1 = 11 - 4x_1 - x_2 - 2x_3 - 10 + 4x_1 + 6x_2 + 2x_3$

$$\Rightarrow w_2 = 1 + 2w_1 + 5x_2$$

Step 2: Improving the solution

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to $w_1 = 5$

$w_2 = 11$

$w_3 = 8$

$$-2 x_1 \quad -3 x_2 \quad - x_3$$

$$-4 x_1 \quad - x_2 \quad -2 x_3$$

$$-3 x_1 \quad -4 x_2 \quad -2 x_3$$

} just rearranging

} what about here?

} “row operations”

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Notice that $w_2 - 2w_1 = 11 - 4x_1 - x_2 - 2x_3 - 10 + 4x_1 + 6x_2 + 2x_3$

$$\Rightarrow w_2 = 1 + 2w_1 + 5x_2$$

x_1 has been eliminated

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

basic variables

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 =$	2.5	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 =$	1	$+2 w_1$	$+5 x_2$	
$w_3 =$	0.5	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

basic variables



nonbasic variables

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 =$	2.5	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 =$	1	$+2 w_1$	$+5 x_2$	
$w_3 =$	0.5	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

basic variables



nonbasic variables

$$w_1 = 0, x_2 = 0, x_3 = 0$$

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 =$	2.5	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 =$	1	$+2 w_1$	$+5 x_2$	
$w_3 =$	0.5	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$



nonbasic variables

basic variables

$$w_1 = 0, x_2 = 0, x_3 = 0 \quad x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

the objective function value has increased

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

basic variables



nonbasic variables

$$w_1 = 0, x_2 = 0, x_3 = 0 \quad x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

The New Dictionary

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

the objective function value has increased

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

basic variables



nonbasic variables

$$w_1 = 0, x_2 = 0, x_3 = 0 \quad x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Which variable should we try to increase next?

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

For w_3 , x_3 can become as large as $0.5/0.5 = 1$.

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_3 = 1, \quad w_1 = x_2 = 0$$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

For w_3 , x_3 can become as large as $0.5/0.5 = 1$.

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_3 = 1, w_1 = x_2 = 0 \qquad x_1 = 2, w_2 = 1, w_3 = 0$$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

For w_3 , x_3 can become as large as $0.5/0.5 = 1$.

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to $x_1 = 2.5 - 0.5 w_1 - 1.5 x_2 - 0.5 x_3$

$w_2 = 1 + 2 w_1 + 5 x_2$

$w_3 = 0.5 + 1.5 w_1 + 0.5 x_2 - 0.5 x_3$

entering variable

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$x_3 = 1, w_1 = x_2 = 0 \qquad x_1 = 2, w_2 = 1, w_3 = 0$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

For w_3 , x_3 can become as large as $0.5/0.5 = 1$.

Step 3: Improving the solution

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to $x_1 = 2.5 - 0.5 w_1 - 1.5 x_2 - 0.5 x_3$

$w_2 = 1 + 2 w_1 + 5 x_2$

$w_3 = 0.5 + 1.5 w_1 + 0.5 x_2 - 0.5 x_3$

entering variable

leaving variable

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$x_3 = 1, w_1 = x_2 = 0 \qquad x_1 = 2, w_2 = 1, w_3 = 0$

For x_1 , x_3 can become as large as $2.5/0.5 = 5$.

For w_2 , x_3 can become as large as ∞ .

For w_3 , x_3 can become as large as $0.5/0.5 = 1$.

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$

$w_2 = 1$

$x_3 = 1$

$-2w_1 - 2x_2 + w_3$

$+2w_1 + 5x_2$

$+3w_1 + x_2 - 2w_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$

$w_2 = 1$

$x_3 = 1$

$-2w_1 - 2x_2 + w_3$

$+2w_1 + 5x_2$

$+3w_1 + x_2 - 2w_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables



nonbasic variables

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$

$w_2 = 1$

$x_3 = 1$

$-2w_1 - 2x_2 + w_3$

$+2w_1 + 5x_2$

$+3w_1 + x_2 - 2w_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables




nonbasic variables

$w_1 = 0, x_2 = 0, w_3 = 0$

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$	 nonbasic variables
$w_2 = 1$	$+2w_1 + 5x_2$	
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$	

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

basic variables

$$w_1 = 0, x_2 = 0, w_3 = 0 \quad x_1 = 2, w_2 = 1, w_3 = 1$$

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

the objective function value has increased

subject to

$$x_1 = 2$$

$$w_2 = 1$$

$$x_3 = 1$$

$$-2w_1 - 2x_2 + w_3$$

$$+2w_1 + 5x_2$$

$$+3w_1 + x_2 - 2w_3$$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$



nonbasic variables

basic variables

$$w_1 = 0, x_2 = 0, w_3 = 0 \quad x_1 = 2, w_2 = 1, w_3 = 1$$

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

the objective function value has increased

subject to

$$x_1 = 2$$

$$w_2 = 1$$

$$x_3 = 1$$

$$-2w_1 - 2x_2 + w_3$$

$$+2w_1 + 5x_2$$

$$+3w_1 + x_2 - 2w_3$$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$



nonbasic variables

basic variables

$$w_1 = 0, x_2 = 0, w_3 = 0 \quad x_1 = 2, w_2 = 1, w_3 = 1$$

Which variable should we try to increase next?

The New Dictionary

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

the objective function value has increased

subject to

$$x_1 = 2$$

$$w_2 = 1$$

$$x_3 = 1$$

$$-2w_1 - 2x_2 + w_3$$

$$+2w_1 + 5x_2$$

$$+3w_1 + x_2 - 2w_3$$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$



nonbasic variables

basic variables

$$w_1 = 0, x_2 = 0, w_3 = 0 \quad x_1 = 2, w_2 = 1, w_3 = 1$$

Which variable should we try to increase next?

We have computed an optimal solution!

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .
2. Write the original dictionary.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

Let's do it again, “mechanically”

Maximise $5x_1 + 4x_2 + 3x_3$

subject to

$$2x_1 + 3x_2 + x_3 \leq 5$$
$$4x_1 + x_2 + 2x_3 \leq 11$$
$$3x_1 + 4x_2 + 2x_3 \leq 8$$
$$x_1, x_2, x_3 \geq 0$$

1. Introduce slack variables

$x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

Maximise $\zeta = 5x_1 + 4x_2 + 3x_3$

subject to

$$w_1 = 5 - 2x_1 + 3x_2 + x_3$$
$$w_2 = 11 - 4x_1 + x_2 + 2x_3 + 3$$
$$w_3 = 8 - 3x_1 + 4x_2 + 2x_3$$
$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

2. Write the original dictionary.

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

Maximise $\zeta =$ $+5 \ x_1$ $+4 \ x_2$ $+3 \ x_3$

subject to $w_1 = 5$ $-2 \ x_1$ $-3 \ x_2$ $- \ x_3$
 $w_2 = 11$ $-4 \ x_1$ $- \ x_2$ $-2 \ x_3$
 $w_3 = 8$ $-3 \ x_1$ $-4 \ x_2$ $-2 \ x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

Maximise $\zeta =$ $+5 \ x_1$ $+4 \ x_2$ $+3 \ x_3$

subject to $w_1 = 5$ $-2 \ x_1$ $-3 \ x_2$ $- \ x_3$
 $w_2 = 11$ $-4 \ x_1$ $- \ x_2$ $-2 \ x_3$
 $w_3 = 8$ $-3 \ x_1$ $-4 \ x_2$ $-2 \ x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 \ x_1$ $+4 \ x_2$ $+3 \ x_3$

subject to $w_1 = 5$ $-2 \ x_1$ $-3 \ x_2$ $- \ x_3$
 $w_2 = 11$ $-4 \ x_1$ $- \ x_2$ $-2 \ x_3$
 $w_3 = 8$ $-3 \ x_1$ $-4 \ x_2$ $-2 \ x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to

$$w_1 = 5$$

$$w_2 = 11$$

$$w_3 = 8$$

$$-2 x_1 - 3 x_2 - x_3$$

$$-4 x_1 - x_2 - 2 x_3$$

$$-3 x_1 - 4 x_2 - 2 x_3$$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

$$5/2 \text{ vs } 11/4 \text{ vs } 8/3 \Rightarrow w_1$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta =$ $+5 x_1$ $+4 x_2$ $+3 x_3$ entering variable

subject to $w_1 = 5$ leaving variable

$w_2 = 11$

$w_3 = 8$

$-2 x_1$	$-3 x_2$	$- x_3$
$-4 x_1$	$- x_2$	$-2 x_3$
$-3 x_1$	$-4 x_2$	$-2 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

$$x_1 = x_2 = x_3 = 0$$

$$w_1 = 5, w_2 = 11, w_3 = 8$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

$$5/2 \text{ vs } 11/4 \text{ vs } 8/3 \Rightarrow w_1$$

5. Increase the value of the entering variable to be $x_k = \min_{i:\hat{a}_{ik}>0} \hat{b}_i/\hat{a}_{ik}$

Maximise $\zeta =$ +5 x_1 +4 x_2 +3 x_3

subject to

$w_1 = 5$	$-2 x_1$	$-3 x_2$	$- x_3$
$w_2 = 11$	$-4 x_1$	$- x_2$	$-2 x_3$
$w_3 = 8$	$-3 x_1$	$-4 x_2$	$-2 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 2.5, x_2 = 0, x_3 = 0$$

6. Compute the new dictionary making sure x_k only appears on the left.

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = x_3 = 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;


4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

entering variable 

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;


4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

entering variable 

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

entering variable

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

$$2.5/0.5 \text{ vs } \infty \text{ vs } 0.5/0.5 \Rightarrow w_3$$

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$

leaving variable \rightarrow w_3

$-0.5 x_3$ \leftarrow entering variable

$$w_1 = x_2 = x_3 = 0$$

$$x_1 = 2.5, w_2 = 1, w_3 = 0.5$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik} x_k \geq 0$).

$$2.5/0.5 \text{ vs } \infty \text{ vs } 0.5/0.5 \Rightarrow w_3$$

5. Increase the value of the entering variable to be $x_k = \min_{i:\hat{a}_{ik}>0} \hat{b}_i/\hat{a}_{ik}$

Maximise $\zeta = 12.5 - 2.5 w_1 - 3.5 x_2 + 0.5 x_3$

subject to

$x_1 = 2.5$	$-0.5 w_1$	$-1.5 x_2$	$-0.5 x_3$
$w_2 = 1$	$+2 w_1$	$+5 x_2$	
$w_3 = 0.5$	$+1.5 w_1$	$+0.5 x_2$	$-0.5 x_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$x_1 = 2.5, x_2 = 0, x_3 = 1$$

6. Compute the new dictionary making
sure x_k only appears on the left.

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = w_3 = 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = w_3 = 0$$

$$x_1 = 2, w_2 = 1, w_3 = 1$$

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

The Simplex Method

1. Introduce slack variables $x_{n+1}, x_{n+2}, \dots, x_m$ and ζ .

2. Write the original dictionary.

Repeat:

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

4. Choose a variable to **enter** the **basis** (**entering variable**) and a variable to **leave** the basis (**leaving variable**).

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

5. Increase the value of the entering variable to be $x_k = \min_{i: \hat{a}_{ik} > 0} \hat{b}_i / \hat{a}_{ik}$

6. Compute the new dictionary making sure x_k only appears on the left.

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = w_3 = 0$$

$$x_1 = 2, w_2 = 1, w_3 = 1$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Maximise $\zeta = 13 - w_1 - 2x_2 - w_3$

subject to

$x_1 = 2$	$-2w_1 - 2x_2 + w_3$
$w_2 = 1$	$+2w_1 + 5x_2$
$x_3 = 1$	$+3w_1 + x_2 - 2w_3$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

$$w_1 = x_2 = w_3 = 0 \qquad x_1 = 2, w_2 = 1, w_3 = 1$$

We have computed an optimal solution!

Potential Problem

Potential Problem

Consider the following LP:

Potential Problem

Consider the following LP:

Maximise $-2x_1 - x_2$

subject to

$$\begin{aligned} -x_1 + x_2 &\leq -1 \\ -x_1 - 2x_2 &\leq -2 \\ x_2 &\leq 1 \\ x_1, x_2 &\geq 0 \end{aligned}$$

Corresponding dictionary

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $w_1 = -1$ $+ \ x_1$ $- \ x_2$
 $w_2 = -2$ $+ \ x_1$ $+2 \ x_2$
 $w_3 = 1$ $- \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

Corresponding dictionary

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $w_1 = -1$ $+ \ x_1$ $- \ x_2$
 $w_2 = -2$ $+ \ x_1$ $+2 \ x_2$
 $w_3 = 1$ $- \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

Corresponding dictionary

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $w_1 = -1$ $+ \ x_1$ $- \ x_2$
 $w_2 = -2$ $+ \ x_1$ $+2 \ x_2$
 $w_3 = 1$ $- \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to **0**.

$$w_1 = x_2 = x_3 = 0$$

Corresponding dictionary

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $w_1 = -1$ $+ \ x_1$ $- \ x_2$
 $w_2 = -2$ $+ \ x_1$ $+2 \ x_2$
 $w_3 = 1$ $- \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = x_2 = x_3 = 0$$

$$w_1 = -1, w_2 = -2, w_3 = 1$$

Corresponding dictionary

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $w_1 = -1$ $+ \ x_1$ $- \ x_2$
 $w_2 = -2$ $+ \ x_1$ $+2 \ x_2$
 $w_3 = 1$ $- \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = x_2 = x_3 = 0 \qquad w_1 = -1, w_2 = -2, w_3 = 1$$

The dictionary is infeasible!

Initialisation

Consider the following LP:

Maximise $-2x_1 - x_2$

subject to

$$\begin{aligned} -x_1 + x_2 &\leq -1 \\ -x_1 - 2x_2 &\leq -2 \\ x_2 &\leq 1 \\ x_1, x_2 &\geq 0 \end{aligned}$$

Initialisation

Consider the following alternative LP:

Maximise $-x_0$

subject to

$$\begin{aligned} -x_1 + x_2 - x_0 &\leq -1 \\ -x_1 - 2x_2 - x_0 &\leq -2 \\ x_2 - x_0 &\leq 1 \\ x_1, x_2, x_0 &\geq 0 \end{aligned}$$

Initialisation

subject to

$$\begin{aligned} -x_1 + x_2 &\leq -1 \\ -x_1 - 2x_2 &\leq -2 \\ x_2 &\leq 1 \\ x_1, x_2 &\geq 0 \end{aligned}$$

Maximise

$$-x_0$$

subject to

$$\begin{aligned} -x_1 + x_2 - x_0 &\leq -1 \\ -x_1 - 2x_2 - x_0 &\leq -2 \\ x_2 - x_0 &\leq 1 \\ x_1, x_2, x_0 &\geq 0 \end{aligned}$$

Initialisation

subject to

$$-x_1 + x_2 \leq -1$$

$$-x_1 - 2x_2 \leq -2$$

$$x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

The first LP is feasible if and only if the second LP has an optimal solution of value 0.

Maximise

$$-x_0$$

subject to

$$-x_1 + x_2 - x_0 \leq -1$$

$$-x_1 - 2x_2 - x_0 \leq -2$$

$$x_2 - x_0 \leq 1$$

$$x_1, x_2, x_0 \geq 0$$

Initialisation

Consider the following alternative LP:

Maximise $-x_0$

subject to

$$\begin{aligned} -x_1 + x_2 - x_0 &\leq -1 \\ -x_1 - 2x_2 - x_0 &\leq -2 \\ x_2 - x_0 &\leq 1 \\ x_1, x_2, x_0 &\geq 0 \end{aligned}$$

Auxiliary problem dictionary

Maximise $\zeta =$ - x_0

$$\begin{array}{llll} \text{subject to} & w_1 = -1 & + x_1 & - x_2 & + x_0 \\ & w_2 = -2 & + x_1 & + 2 x_2 & + x_0 \\ & w_3 = 1 & & - x_2 & + x_0 \end{array}$$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

Auxiliary problem dictionary

Maximise $\zeta =$ $-x_0$

subject to

$w_1 = -1$	$+ x_1$	$- x_2$	$+ x_0$
$w_2 = -2$	$+ x_1$	$+2 x_2$	$+ x_0$
$w_3 = 1$		$- x_2$	$+ x_0$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

Auxiliary problem dictionary

Maximise $\zeta =$ $-x_0$

subject to

$w_1 = -1$	$+ x_1$	$- x_2$	$+ x_0$
$w_2 = -2$	$+ x_1$	$+2 x_2$	$+ x_0$
$w_3 = 1$		$- x_2$	$+ x_0$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

The dictionary is infeasible!

Auxiliary problem dictionary

Maximise $\zeta =$ $-x_0$

subject to

$w_1 = -1$	$+ x_1$	$- x_2$	$+ x_0$
$w_2 = -2$	$+ x_1$	$+2 x_2$	$+ x_0$
$w_3 = 1$		$- x_2$	$+ x_0$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

The dictionary is infeasible!

Entering variable: x_0

Auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to

$w_1 = -1$	$+ x_1$	$- x_2$	$+ x_0$
$w_2 = -2$	$+ x_1$	$+2 x_2$	$+ x_0$
$w_3 = 1$		$- x_2$	$+ x_0$

entering variable

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

The dictionary is infeasible!

Entering variable: x_0

Auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to

$w_1 = -1$	$+ x_1$	$- x_2$	$+ x_0$
$w_2 = -2$	$+ x_1$	$+2 x_2$	$+ x_0$
$w_3 = 1$		$- x_2$	$+ x_0$

entering variable

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

The dictionary is infeasible!

Entering variable: x_0

Leaving variable: the one that is “most infeasible”

Auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to $w_1 = -1$
 $w_2 = -2$
 $w_3 = 1$

leaving variable



$+ x_1$	$- x_2$	$+ x_0$
$+ x_1$	$+2 x_2$	$+ x_0$
	$- x_2$	$+ x_0$

entering variable



$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

The dictionary is infeasible!

Entering variable: x_0

Leaving variable: the one that is “most infeasible”

Auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to $w_1 = -1$
 $w_2 = -2$
 $w_3 = 1$

leaving variable



$+ x_1$	$- x_2$	$+ x_0$
$+ x_1$	$+2 x_2$	$+ x_0$
	$- x_2$	$+ x_0$

entering variable



$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

3. Find a basic feasible solution by setting the nonbasic variables to 0.

The dictionary is infeasible!

Entering variable: x_0

Leaving variable: the one that is “most infeasible”

6. Compute the new dictionary making sure x_0 only appears on the left.

The new auxiliary problem dictionary

Maximise $\zeta = -2 \quad + \quad x_1 \quad + 2 \quad x_2 \quad - \quad w_2$

subject to

$w_1 =$	1		$-3 \quad x_2$	$+ \quad w_2$
$x_0 =$	2	$- \quad x_1$	$-2 \quad x_2$	$+ \quad w_2$
$w_3 =$	3	$- \quad x_1$	$-3 \quad x_2$	$+ \quad w_2$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

The new auxiliary problem dictionary

Maximise $\zeta = -2 + x_1 + 2x_2 - w_2$

subject to

$w_1 =$	1		$-3x_2$	$+ w_2$
$x_0 =$	2	$-x_1$	$-2x_2$	$+ w_2$
$w_3 =$	3	$-x_1$	$-3x_2$	$+ w_2$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

The dictionary is feasible, we can apply the simplex method.

The new auxiliary problem dictionary


Maximise $\zeta = -2 + x_1 + 2x_2 - w_2$

subject to

$w_1 = 1$		$-3x_2$	$+ w_2$
$x_0 = 2$	$-x_1$	$-2x_2$	$+ w_2$
$w_3 = 3$	$-x_1$	$-3x_2$	$+ w_2$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

The dictionary is feasible, we can apply the simplex method.



steps...

The final auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to

$x_2 = 0.33$		$-0.33 w_1$	$+0.33 w_2$
$x_1 = 1.33$	$-x_0$	$+0.67 w_1$	$+0.33 w_2$
$w_3 = 2$	$+x_0$	$+0.33 w_1$	$+0.33 w_2$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

The final auxiliary problem dictionary

Maximise $\zeta = -x_0$

subject to

$x_2 = 0.33$		$-0.33 w_1$	$+0.33 w_2$
$x_1 = 1.33$	$-x_0$	$+0.67 w_1$	$+0.33 w_2$
$w_3 = 2$	$+x_0$	$+0.33 w_1$	$+0.33 w_2$

$$x_1, x_2, w_1, w_2, w_3, x_0 \geq 0$$

Remove x_0 from the constraints and substitute the original objective function.

The first dictionary of our original problem

Maximise $\zeta =$ $-2 \ x_1$ $-x_2$

subject to $x_2 =$ 0.33 $-0.33 \ w_1$ $+0.33 \ w_2$
 $x_1 =$ 1.33 $+0.67 \ w_1$ $+0.33 \ w_2$
 $w_3 =$ 2 $+0.33 \ w_1$ $+0.33 \ w_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

The first dictionary of our original problem

Maximise $\zeta =$ $-2 x_1$ $-x_2$

subject to $x_2 =$ 0.33 $-0.33 w_1$ $+0.33 w_2$
 $x_1 =$ 1.33 $+0.67 w_1$ $+0.33 w_2$
 $w_3 =$ 2 $+0.33 w_1$ $+0.33 w_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

We should have only nonbasic variables in the objective function.

Easy Fix

Maximise $\zeta = \quad -2 \ x_1 \quad -x_2$

subject to $w_1 = -1$ $+ \ x_1 \quad - \ x_2$
 $w_2 = -2$ $+ \ x_1 \quad +2 \ x_2$
 $w_3 = \ 1$ $\quad \quad - \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

Easy Fix

Maximise $\zeta = \quad -2 \ x_1 \quad -x_2$

subject to $w_1 = -1 \quad + \ x_1 \quad - \ x_2$
 $w_2 = -2 \quad + \ x_1 \quad +2 \ x_2$
 $w_3 = \ 1 \quad \quad \quad - \ x_2$

$$x_1, x_2, w_1, w_2, w_3 \geq 0$$

We have $\zeta = -2x_1 - x_2 = -3 - w_1 - w_2$

The first dictionary of our original problem

Maximise $\zeta =$ $-3 \ w_1 \quad - \quad w_2$

subject to $x_2 =$ 0.33 $-0.33 \ w_1 \quad +0.33 \ w_2$
 $x_1 =$ 1.33 $+0.67 \ w_1 \quad +0.33 \ w_2$
 $w_3 =$ 2 $+0.33 \ w_1 \quad +0.33 \ w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to

$x_2 = 0.33$	$-0.33 w_1 + 0.33 w_2$
$x_1 = 1.33$	$+0.67 w_1 + 0.33 w_2$
$w_3 = 2$	$+0.33 w_1 + 0.33 w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to

$x_2 = 0.33$	$-0.33 w_1 + 0.33 w_2$
$x_1 = 1.33$	$+0.67 w_1 + 0.33 w_2$
$w_3 = 2$	$+0.33 w_1 + 0.33 w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = w_2 = 0$$

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to

$x_2 = 0.33$	$-0.33 w_1 + 0.33 w_2$
$x_1 = 1.33$	$+0.67 w_1 + 0.33 w_2$
$w_3 = 2$	$+0.33 w_1 + 0.33 w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = w_2 = 0$$

$$x_1 = 1.33, x_2 = 0.33, w_3 = 2$$

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to

$x_2 = 0.33$	$-0.33 w_1 + 0.33 w_2$
$x_1 = 1.33$	$+0.67 w_1 + 0.33 w_2$
$w_3 = 2$	$+0.33 w_1 + 0.33 w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = w_2 = 0 \qquad x_1 = 1.33, x_2 = 0.33, w_3 = 2$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to $x_2 = 0.33$

$x_1 = 1.33$

$w_3 = 2$

$$-0.33 w_1 + 0.33 w_2$$

$$+0.67 w_1 + 0.33 w_2$$

$$+0.33 w_1 + 0.33 w_2$$

We have found an optimal solution!

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = w_2 = 0$$

$$x_1 = 1.33, x_2 = 0.33, w_3 = 2$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

The first dictionary of our original problem

Maximise $\zeta = -3 w_1 - w_2$

subject to

$x_2 = 0.33$	$-0.33 w_1 + 0.33 w_2$
$x_1 = 1.33$	$+0.67 w_1 + 0.33 w_2$
$w_3 = 2$	$+0.33 w_1 + 0.33 w_2$

$$x_1, x_2, x_3, w_1, w_2, w_3 \geq 0$$

We have found an optimal solution!

We were lucky: we can only expect to find a feasible solution.

3. Find a **basic feasible solution** by setting the **nonbasic variables** to 0.

$$w_1 = w_2 = 0$$

$$x_1 = 1.33, x_2 = 0.33, w_3 = 2$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

What if we have this dictionary?

Maximise $\zeta = 5 \quad + \quad x_3 \quad - \quad x_1$

subject to $x_2 = 5$ $+2 \quad x_3 \quad -3 \quad x_1$
 $x_4 = 7$ $-4 \quad x_1$
 $x_5 =$ x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

What if we have this dictionary?

Maximise $\zeta = 5 + x_3 - x_1$

subject to $x_2 = 5$ $+2x_3 - 3x_1$
 $x_4 = 7$ $-4x_1$
 $x_5 =$ x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break;**

What if we have this dictionary?

Maximise $\zeta = 5 + x_3 - x_1$

subject to $x_2 = 5$ $+2x_3 - 3x_1$
 $x_4 = 7$ $-4x_1$
 $x_5 =$ x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	+2	x_3	-3	x_1
$x_4 = 7$			-4	x_1
$x_5 =$				x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

Entering variable: Any variable with positive coefficient in the objective function. If none exists, **break**;

Leaving variable: The variable with the **smallest ratio** \hat{b}_i / \hat{a}_{ik} (for the constraint $\hat{b}_i - \hat{a}_{ik}x_k \geq 0$).

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ entering variable

subject to $x_2 = 5$ $+2 \ x_3 \ -3 \ x_1$
 $x_4 = 7$ $-4 \ x_1$
 $x_5 =$ x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	$+2$	x_3	-3	x_1
$x_4 = 7$			-4	x_1
$x_5 =$				x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	+2	x_3	-3	x_1
$x_4 = 7$			-4	x_1
$x_5 =$				x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

We should not violate any constraints though!

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	$+2 \ x_3 \ -3 \ x_1$
$x_4 = 7$	$ \ -4 \ x_1$
$x_5 =$	$ x_1$

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	+2	x_3	-3	x_1
$x_4 = 7$			-4	x_1
$x_5 =$				x_1

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	$+2 \quad x_3 \quad -3 \quad x_1$
$x_4 = 7$	$ -4 \quad x_1$
$x_5 =$	$ x_1$

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

This does not happen regardless of how much we increase x_3 .

What if we have this dictionary?

Maximise $\zeta = 5 + \boxed{x_3} - x_1$ ← entering variable

subject to

$x_2 = 5$	$+2 \quad x_3 \quad -3 \quad x_1$
$x_4 = 7$	$ -4 \quad x_1$
$x_5 =$	$ x_1$

The LP is unbounded!

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

We can increase the value of some nonbasic variable, here x_3

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

This does not happen regardless of how much we increase x_3 .

What about this dictionary?

Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$


subject to $x_3 = 1 - 0.5 x_1 - 0.5 w_1$
 $w_2 = x_1 - x_2 + w_1$

$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

What about this dictionary?

Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$

subject to $x_3 = 1$ $-0.5 x_1$ $-0.5 w_1$
 $w_2 =$ $x_1 - x_2 + w_1$ **entering variable**



$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

What about this dictionary?

Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$

subject to $x_3 = 1$

leaving variable $\rightarrow w_2 =$

$$\begin{array}{rcl} -0.5 x_1 & & -0.5 w_1 \\ x_1 & - & x_2 & + w_1 \end{array}$$

entering variable


$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

What about this dictionary?

Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$

subject to $x_3 = 1$

$w_2 =$  leaving variable

$$\begin{array}{rcl} -0.5 x_1 & & -0.5 w_1 \\ x_1 & - & x_2 & + w_1 \end{array}$$


entering variable

$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

We can increase the value of some nonbasic variable, here x_2

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

What about this dictionary?



Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$

subject to $x_3 = 1$

$w_2 =$

leaving variable 

$$\begin{array}{rcl} -0.5 x_1 & & -0.5 w_1 \\ x_1 & - & x_2 & + w_1 \end{array}$$

entering variable

$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

We can increase the value of some nonbasic variable, here x_2

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

x_2 cannot be increased! Are we stuck?

What about this dictionary?

Maximise $\zeta = 3 - 0.5 x_1 + 2 x_2 - 1.5 w_1$

subject to $x_3 = 1$

leaving variable $\rightarrow w_2 =$

$$\begin{array}{rcl} -0.5 x_1 & & -0.5 w_1 \\ x_1 & - & x_2 & + w_1 \end{array}$$

entering variable

$$x_1, x_2, x_3, w_1, w_2 \geq 0$$

We can increase the value of some nonbasic variable, here x_2

We should not violate any constraints though!

We don't want any of the slack variables to become negative.

x_2 cannot be increased! Are we stuck?

Degeneracy! Next lecture

Historic Note

The Simplex Method was invented by George Dantzig in 1947.

It is still being used today in most of the LP-solvers.

Historic Note

The Simplex Method was invented by George Dantzig in 1947.

It is still being used today in most of the LP-solvers.

The origins of the simplex method go back to one of two famous unsolved problems in mathematical statistics proposed by Jerzy Neyman, which I mistakenly solved as a homework problem; it later

Dantzig. Origins of the Simplex Method. In *A History of Scientific Computing*, 1990.