

This homework runs from Thursday 3 October 2024 until 12 noon on Thursday 10 October 2023. Submission is to Gradescope Homework 3.

Questions marked with an asterisk * may be a little harder than others. All are still within the course curriculum, though, and can be done using the methods taught in the study guides and textbook.

You should aim to write out solutions that someone who does not already know the answer could follow and understand.

Please remember the good scholarly practice requirements of the University regarding work for credit. You can find guidance at the School page <https://web.inf.ed.ac.uk/infweb/admin/policies/academic-misconduct>. This also has links to the relevant University pages.

Question 1

Prove using the element method that for any four sets A , B , C , and D , if $A \subseteq C$ and $B \subseteq D$ then $(A \cap B) \subseteq (C \cap D)$ and $(A \cup B) \subseteq (C \cup D)$.

[4 marks]

Question 2

Here are three functions from $\mathbb{R}$ to $\mathbb{R}$.

- (a) $f(x) = -x$.
- (b) $g(x) = 2^x$.
- (c) $h(x) = (x^3 - x)$.

For each of these sketch a graph of the function, say whether it is injective, and say whether it is surjective.

[3 marks]

* Question 3

Call a function $s : X \rightarrow Y$ between two sets X and Y a *section* if there is another function (known as a *retraction*) $r : Y \rightarrow X$ such that $r \circ s = I_X$. Here I_X is the identity function on X . Suppose that $f : A \rightarrow B$ and $g : A \rightarrow C$ are both sections.

- (a) Is there necessarily a section $h : A \rightarrow B \cup C$?
- (b) Is there necessarily a section $j : A \rightarrow B \cap C$?
- (c) Is there necessarily a section $k : A \rightarrow B \times C$?

In each case either show how to construct such a section, together with a suitable corresponding retraction, or give a counterexample.

[3 marks]