


Discrete Mathematics and Probability

Week 7



Heather Yorston

Topics

- ▶ Recap: examples with equally likely outcomes
- ▶ Conditional probability: how knowledge influences probability
- ▶ Bayes' theorem: link probabilities of related events

Recap


Permutations and combinations

Example

An urn contains 6 red balls and 5 blue balls.

Draw three balls at random, at once (that is, without replacement).

What is the chance of drawing one red and two blue balls?



Sample space

$$|\Omega| = {}_{11}P_3 = \frac{11!}{(11-3)!} = 11 \times 10 \times 9$$

Event space

$$E = \left\{ \underset{\substack{| \\ 6 \cdot 5 \cdot 4}}{R, B_1, B_2}, \underset{\substack{| \\ 5 \cdot 6 \cdot 4}}{B_1, R, B_2}, \underset{\substack{| \\ 5 \cdot 4 \cdot 6}}{B_1, B_2, R_1} \right\}$$

ways

$$|E| = 3 \times 4 \times 5 \times 6$$

$$P(E) = \frac{|E|}{|\Omega|} = (3) \frac{4 \cdot 5 \cdot 6}{9 \cdot 10 \cdot 11} = \frac{4}{11}$$

OR

$$\begin{aligned} & P(R, B_1, B_2) + P(B_1, R, B_2) + P(B_1, B_2, R_1) \\ &= \frac{6}{11} \times \frac{5}{10} \times \frac{4}{9} + \frac{5}{11} \times \frac{6}{10} \times \frac{4}{9} + \frac{5}{11} \times \frac{4}{10} \times \frac{6}{9} \\ &= 3 \times \left(\frac{6 \cdot 5 \cdot 4}{11 \cdot 10 \cdot 9} \right) \end{aligned}$$

Permutations and combinations

Example

An urn contains n balls, one of which is red, all others are black. We draw k balls at random (without replacement). What is the chance that the red ball will be drawn?

$$S = \{k\text{-combinations of } n\}$$

$$|S| = \binom{n}{k} \quad E = \{\text{pick red ball and } k-1 \text{ others}\}$$

$$|E| = \binom{1}{1} \binom{n-1}{k-1}$$

$$P(E) = \frac{|E|}{|S|} = \frac{\binom{1}{1} \binom{n-1}{k-1}}{\binom{n}{k}} = \frac{\cancel{(n-1)!} \cdot \overset{k}{k!} \cdot \cancel{(n-k)!}}{\cancel{(k-1)!} \cdot \cancel{(n-1-k+1)!} \cdot \overset{n}{n!}} = \frac{k}{n}$$

Alternative Method

Drawing in order:

$$E_i = \{i^{\text{th}} \text{ draw is red}\}.$$


$$P(E_i) = \frac{1}{n}$$

$$E = \cup E_i$$

$$P(E) = P(\cup E_i) = \sum_i P(E_i) = \sum_{i=1}^k \frac{1}{n} = \frac{1}{n} \sum_{i=1}^n 1 = \frac{k}{n}$$

Equally likely events

pigeonhole principle



Example

Out of n people, what is the probability that no two share a birthday?

0 if $n > 365$ (pigeonhole principle)

$\Omega = \{ \text{choice of day for each of } n \text{ people} \}$
[with replacement]

$$|\Omega| = 365^n$$

$E = \{ \text{no coinciding birthdays} \}$

$$|E| = 365 \times 364 \times \dots \times (365 - n + 1)$$

$$P(E) = \frac{|E|}{|\Omega|} = \frac{365!}{(365-n)! 365^n} =$$

$$= \frac{365!}{(365-n)!}$$

n	$P(E)$	
10	88%	50/37%
20	59%	
30	29%	
40	11%	

Conditional probability

Conditional probability

Often you have *partial information* about the outcome of an experiment. This alters the likelihoods for various outcomes.

Example

Roll two dice. What is the probability that the sum of the numbers is 8? What if we know that the first die shows a 5?

$$\Omega = \{1, \dots, 6\}^2$$
$$E = \{(2,6), (3,5), \dots, (6,2)\} \quad P(E) = \frac{5}{36}$$

but if first die is 5 now:

$$\Omega = \{1, 2, \dots, 6\}$$

$$F = \{3\}$$

$$P(F) = \frac{1}{6}$$

partial info
changes prob'.

Reduced sample space

We reduced our world to the event we were given:

$$F = \{\text{first die shows 5}\} = \{(5, 1), (5, 2), \dots, (5, 6)\}$$

Definition

The event that is given to us is called a *reduced sample space*. We can simply work in this set to figure out the conditional probabilities given this event.

The event F has 6 equally likely outcomes. Only one of them, $(5, 3)$, provides a sum of 8. Hence the conditional probability is $\frac{1}{6}$.

Definition of conditional probability

The question can be reformulated.

$$E = \{\text{the sum is 8}\} = \{(2, 6), (3, 5), \dots, (6, 2)\}$$

“In what proportion of cases in F will E also occur?”

“How does probability of ‘ E and F ’ compare to probability of F ?”

Definition

Let F be an event with $\mathbf{P}(F) > 0$.

The *conditional probability of E given F* is:

$$\mathbf{P}(E|F) = \frac{\mathbf{P}(E \cap F)}{\mathbf{P}(F)}$$

A handwritten diagram shows a Venn diagram with two overlapping circles. The left circle is labeled $P(\bar{F})$ and the right circle is labeled $P(F)$. The intersection of the two circles is labeled $P(E|F)$. To the right of the diagram, the following equations are written:

$$P(E \cap F) = P(F) P(E|F)$$
$$\text{so } P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Axioms

Proposition

Conditional probability $\mathbf{P}(\cdot | F)$ satisfies the axioms of probability:

1. conditional probability is non-negative: $\mathbf{P}(E | F) \geq 0$;
2. conditional probability of sample space is one: $\mathbf{P}(\Omega | F) = 1$;
3. for countably many *mutually exclusive* events E_1, E_2, \dots :

$$\mathbf{P}\left(\bigcup_i E_i \mid F\right) = \sum_i \mathbf{P}(E_i \mid F)$$

How to compute conditional probabilities

Corollary

- ▶ $\mathbf{P}(E^c | F) = 1 - \mathbf{P}(E | F)$
- ▶ $\mathbf{P}(\emptyset | F) = 0$
- ▶ $\mathbf{P}(E | F) = 1 - \mathbf{P}(E^c | F) \leq 1$
- ▶ $\mathbf{P}(E \cup G | F) = \mathbf{P}(E | F) + \mathbf{P}(G | F) - \mathbf{P}(E \cap G | F)$
- ▶ If $E \subseteq G$, then $\mathbf{P}(G - E | F) = \mathbf{P}(G | F) - \mathbf{P}(E | F)$
- ▶ If $E \subseteq G$, then $\mathbf{P}(E | F) \leq \mathbf{P}(G | F)$

BUT: Don't change the condition!

$\mathbf{P}\{E | F\}$ and $\mathbf{P}\{E | F^c\}$ have nothing to do with each other.

Multiplication rule

Proposition (*Multiplication rule*)

$$\mathbf{P}(E_1 \cap \cdots \cap E_n) = \mathbf{P}(E_1) \cdot \mathbf{P}(E_2 | E_1) \cdot \mathbf{P}(E_3 | E_1 \cap E_2) \\ \cdots \mathbf{P}(E_n | E_1 \cap \cdots \cap E_{n-1})$$

Example again

Example

An urn contains 6 red and 5 blue balls. We draw three balls at random, at once (that is, without replacement). What is the chance of drawing one red and two blue balls?

$$\begin{aligned} & p(R_1 \cap B_1 \cap B_2) + p(B_1 \cap R_1 \cap R_2) + \\ & \quad p(B_1 \cap B_2 \cap R_1) \\ & = p(R_1) \cdot p(B_1 | R_1) \cdot p(B_2 | (R_1 \cap B_1)) + \dots \end{aligned}$$

Bayes' theorem

Bayes' Theorem

The aim is to say something about $P(F | E)$, once we know $P(E | F)$ (and other things. . .). This will be very useful, and serve as a fundamental tool in probability and statistics.

The Law of Total Probability

Theorem (*Partition Theorem*)

$$\mathbf{P}(E) = \mathbf{P}(E | F) \cdot \mathbf{P}(F) + \mathbf{P}(E | F^c) \cdot \mathbf{P}(F^c)$$

The Law of Total Probability

Theorem (*Partition Theorem*)

$$\mathbf{P}(E) = \mathbf{P}(E | F) \cdot \mathbf{P}(F) + \mathbf{P}(E | F^c) \cdot \mathbf{P}(F^c)$$

Definition

Countably many events F_1, F_2, \dots form a *partition* of Ω if $F_i \cap F_j = \emptyset$ and $\bigcup_i F_i = \Omega$.

Theorem (*Partition Theorem*)

For any event E and any partition F_1, F_2, \dots :

$$\mathbf{P}(E) = \sum_i \mathbf{P}(E | F_i) \cdot \mathbf{P}(F_i)$$

Summary

- ▶ Probability: multiple ways to compute
- ▶ Conditional probability: reduced sample space, multiplication rule
- ▶ Bayes' theorem: partition theorem, belief update