



THE UNIVERSITY
of EDINBURGH

Methods for Causal Inference

Lecture 11: Front-Door Criterion

Ava Khamseh

School of Informatics
2025-2026


The adjustment formula

T: Drug usage

X: Sex

Y: Recovery

Use P_m as an
intermediate tool



To know how effective the drugs is in the population, compare the **hypothetical interventions** by which

- (i) the drug is administered uniformly to the entire population $do(T=1)$ vs
- (ii) complement, i.e., everyone is prevented from taking the drug $do(T=0)$

Aim: Estimate the difference (**Average Causal Effect ACE, aka ATE**)

$$p(Y = 1|do(T = 1)) - p(Y = 1|do(T = 0))$$

The Backdoor Criterion

Under what conditions does a causal model permit computing the causal effect of one variable on another, from **data** obtained from **passive observations**, with **no intervention**? i.e.,



Under what conditions is the structure of a causal graph sufficient of computing a causal effect from a given data set? **Identifiability**

Backdoor Criterion: Given an ordered pair of variables (T,Y) in a DAG G, a set of variables X satisfies the backdoor criterion relative to (T,Y) if:

- (i) no node in X is a descendent of T
- (ii) X block every path between T and Y that contains an arrow into T

If X satisfies the backdoor criterion then the causal effect of T on Y is given by:

$$p(Y = y|do(T = t)) = \sum_x p(Y = y|T = t, X = x)p(X = x)$$



The Backdoor Criterion

Under what conditions does a causal model permit computing the causal effect of one variable on another, from **data** obtained from **passive observations**, with **no intervention**? i.e.,

Under what conditions is the structure of a causal graph sufficient of computing a causal effect from a given data set? Identifiability

Backdoor Criterion: Given an ordered pair of variables (T,Y) in a DAG G, a set of variables X satisfies the backdoor criterion relative to (T,Y) if:

- (i) no node in X is a descendent of T
- (ii) X block every path between T and Y that contains an arrow into T

If X satisfies the backdoor criterion then the causal effect of T on Y is given by:


$$p(Y = y|do(T = t)) = \sum_x p(Y = y|T = t, X = x)p(X = x)$$

In other words, condition on a set of nodes X such that:

- (i) We block all spurious paths between T and Y
- (ii) We leave all direct paths from T to Y unperturbed
- (iii) We create no new spurious paths (do not unblock any new paths)

Recall ...

- Backdoor does not exhaust all ways of estimating causal effects from a graph
- Front-door criterion can still be used for patterns that do not satisfy the backdoor criterion
- Example: Smoking and lung cancer (1970), industry argued to prevent antismoking regulation by suggesting that the correlation could be explained by a carcinogenic genotype that induces a craving for nicotine
- Recall sensitivity analysis
- Recall instrumental variable approach



Instrumental Variable assumptions

- **SUTVA:** Potential outcomes for each individual i are unrelated to the treatment status of other individuals:

$$Y^{(i)}(\mathbf{Z}, \mathbf{T}) = Y^{(i)}(Z^{(i)}, T^{(i)}) , \quad |\mathbf{Z}| = |\mathbf{T}| = N \text{ individuals}$$


- **Non-zero average/relevant:** Treatment assignment Z associated with the treatment $\mathbb{E} \left[\left(T^{(i)} | z = 1 \right) - \left(T^{(i)} | z = 0 \right) \right]$
- Treatment assignment Z is random (Z and Y do not share a cause).

$$\left(Y^{(i)} | z = 1, t \right) = \left(Y^{(i)} | z = 0, t \right)$$

- **Exclusion Restriction:** Any effect of Z on Y is via an effect of Z on T , i.e., Z should not affect Y when T is held constant


- **Monotonicity** (increasing encouragement “dose” increases probability of treatment, no defiers):

$$\left(T^{(i)} | z = 1 \right) \geq \left(T^{(i)} | z = 0 \right)$$



Overview of the course

- **Lecture 1:** Introduction & Motivation, why do we care about causality? Why deriving causality from observational data is non-trivial.
- **Lecture 2:** Recap of probability theory, variables, events, conditional probabilities, independence, law of total probability, Bayes' rule
- **Lecture 3:** Recap of regression, multiple regression, graphs, SCM
- **Lecture 4-20:**



Pearl's Front-Door Criterion: An Example

- Fig (a): The graph does not satisfy the backdoor, since the quantity we need to condition on to block the path, i.e. the genotype, is unobserved
- Fig (b): Additional measurement available: tar deposits in patients lungs
- Fig (b) still does not satisfy the backdoor criterion but we can determine the causal effect:

$$p(Y = y | do(X = x))$$




Figure 3.10 A graphical model representing the relationships between smoking (X) and lung cancer (Y), with unobserved confounder (U) and a mediating variable Z

Pearl's Front-Door Criterion: A crafted example

Interpretation 1: Tobacco industry

Beneficial effect of smoking:

15% of smokers have developed lung cancer vs 90.25% of non-smokers **within tar and non-tar** subgroups, smokers have a much lower percentage of cancer than non-smokers (numbers in the table are engineered to illustrate the point that observations are not to be trusted)

Table 3.1 A hypothetical data set of randomly selected samples showing the percentage of cancer cases for smokers and nonsmokers in each tar category (numbers in thousands)

	Tar 400		No tar 400		All subjects 800	
	Smokers	Nonsmokers	Smokers	Nonsmokers	Smokers	Nonsmokers
No cancer	380 323 (85%)	20 1 (5%)	20 18 (90%)	380 38 (10%)	400 341 (85%)	400 39 (9.75%)
Cancer	57 (15%)	19 (95%)	2 (10%)	342 (90%)	59 (15%)	361 (90.25%)

Pearl's Front-Door Criterion: A crafted example

Interpretation 2: Anti-smoking lobbyists

Smoking **increases** the risk of lung cancer

If one chooses to smoke, then one's chances of building tar deposits are 95% (380/400) vs 5% (20/400) for the non-cancer group.

To evaluate effect of tar, look at **smokers and non-smokers separately**. Tar has harmful effects in both groups: in smokers it increases risk of cancer from 10% to 15% and in non-smokers 90% to 95%. Therefore: Smoking -> tar -> cancer.

Regardless of any natural craving, avoid harmful tar by not smoking.

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

Pearl's Front-Door Criterion

$X \rightarrow Z$ is identifiable, since no back path from X and Z : $X \leftarrow U \rightarrow Y \leftarrow Z$


$$p(Z = z | do(X = x)) = p(Z = z | X = x) \quad *$$

$Z \rightarrow Y$ is identifiable, since backdoor from Z to Y :

$$Z \leftarrow X \leftarrow U \rightarrow Y$$

is blocked by conditioning on X :

$$p(Y = y | do(Z = z)) = \sum_x p(Y = y | Z = z, X = x) p(X = x) \quad **$$



Pearl's Front-Door Criterion

Letting z be the value Z takes when setting $X=x$, from the graph, we have:

$$p(Y | do(X = x)) = p(Y | do(X = x), Z) = p(Y | do(Z = z))$$

Then summing over all states z of Z :


$$p(Y = y | do(X = x)) = \sum_z p(Y = y, z | do(X = x)) \quad \text{Total prob rule}$$

Product rule:

$$= \sum_z p(Y = y | z, do(X = x)) p(z | do(X = x))$$

Line 1

$$= \sum_z p(Y = y | do(Z = z)) p(z | do(X = x))$$



Pearl's Front-Door Criterion


$$p(Z = z | do(X = x)) = p(Z = z | X = x) \quad *$$

$$p(Y = y | do(Z = z)) = \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') \quad **$$

$$p(Y = y | do(X = x)) = \sum_z p(Y = y | do(Z = z)) p(Z = z | do(X = x))$$

Using * and ** summing over all states z of Z :

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$



Front-door formula

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations for the above equation:

- 342/380 points to $p(Y = 1 | z = 0, x' = 0)$
- 2/20 points to $p(x' = 0)$
- 19/20 points to $p(z = 0 | x = 1)$
- 57/380 points to $p(Y = 1 | z = 1, x' = 1)$
- 0.5 points to $p(x' = 1)$
- 20/400 points to $p(z = 1 | x = 1)$
- 380/400 points to $p(z = 0 | x = 1)$

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations: $342/380$ points to the first term; $2/20$ points to the second term; $19/20$ points to the third term; $57/380$ points to the fourth term. 0.5 points to the boxed $p(x' = 1)$ term. $20/400$ points to the $p(z = 1 | x = 1)$ term. $380/400$ points to the $p(z = 0 | x = 1)$ term.

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations: $342/380$ points to the first term; $2/20$ points to the second term; $19/20$ points to the third term; $57/380$ points to the fourth term. A red box highlights the conditional probabilities $p(z = 0 | x = 1)$ and $p(z = 1 | x = 1)$. Arrows indicate $p(z = 0 | x = 1) = 0.5$ and $p(z = 1 | x = 1) = 380/400$.

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations:
 - $342/380$ points to $p(Y = 1 | z = 0, x' = 0)$
 - $2/20$ points to $p(x' = 0) p(z = 0 | x = 1)$
 - $19/20$ points to $p(x' = 1) p(z = 0 | x = 1)$
 - $57/380$ points to $p(Y = 1 | z = 1, x' = 0)$
 - 0.5 points to $p(x' = 1) p(z = 1 | x = 1)$
 - $20/400$ points to $p(z = 1 | x = 1)$
 - $380/400$ points to $p(z = 0 | x = 1)$

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (90%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations:
 - $342/380$ points to $p(Y = 1 | z = 0, x' = 0)$
 - $2/20$ points to $p(Y = 1 | z = 0, x' = 1)$
 - $19/20$ points to $p(Y = 1 | z = 1, x' = 0)$
 - $57/380$ points to $p(Y = 1 | z = 1, x' = 1)$
 - 0.5 points to $p(x' = 1)$
 - $20/400$ points to $p(z = 0 | x = 1)$
 - $380/400$ points to $p(z = 1 | x = 1)$

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ \boxed{p(Y = 1 | z = 1, x' = 0)} p(x' = 0) p(z = 1 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1) \\
 &= 0.5475
 \end{aligned}$$

Annotations: 342/380, 2/20, 19/20, 57/380, 20/400, 0.5, 380/400

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase

Pearl's Front-Door Criterion: Which group is right?

$$p(Y = y | do(X = x)) = \sum_z \sum_{x'} p(Y = y | Z = z, X = x') p(X = x') p(Z = z | X = x)$$

$$\begin{aligned}
 p(Y = 1 | do(X = 1)) &= p(Y = 1 | z = 0, x' = 0) p(x' = 0) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 0, x' = 1) p(x' = 1) p(z = 0 | x = 1) \\
 &+ p(Y = 1 | z = 1, x' = 0) p(x' = 0) p(z = 1 | x = 1) \\
 &+ \boxed{p(Y = 1 | z = 1, x' = 1) p(x' = 1) p(z = 1 | x = 1)} \\
 &= 0.5475
 \end{aligned}$$

Annotations: $342/380$ (points to $p(Y=1|z=0, x'=0)$), $2/20$ (points to $p(x'=1)$), $19/20$ (points to $p(Y=1|z=1, x'=0)$), $57/380$ (points to $p(Y=1|z=1, x'=1)$), 0.5 (points to $p(x'=1)$), $20/400$ (points to $p(z=0|x=1)$), $380/400$ (points to $p(z=1|x=1)$)

$$p(Y = 1 | do(X = 0)) = 0.5025$$

Average Causal Effect ACE: $p(Y = 1 | do(X = 1)) - p(Y = 1 | do(X = 0)) = 0.045$

Table 3.2 Reorganization of the data set of Table 3.1 showing the percentage of cancer cases in each smoking-tar category (numbers in thousands)

	Smokers 400		Nonsmokers 400		All subjects 800	
	Tar	No tar	Tar	No tar	Tar	No tar
No cancer	380 (85%)	20 (90%)	20 (5%)	380 (10%)	400 (81%)	400 (19%)
Cancer	57 (15%)	2 (10%)	19 (95%)	342 (90%)	76 (19%)	344 (81%)

4.5% increase


Pearl's Front-Door Adjustment

Front-door criterion: A set of variables Z is said to satisfy the front-door criterion relative to (X,Y) if:

1. Z intercepts all directed paths from X to Y
2. There is no unblocked path from X to Z
3. All backdoor paths from Z to Y are blocked by X

Front-door adjustment: If Z satisfied the front-door criterion relative to (X,Y) , and if $p(x,z) > 0$, then the causal effect of X on Y is identifiable and is given by:

$$p(y|do(x)) = \sum_z p(z|x) \sum_{x'} p(y|x', z) p(x')$$



Pearl's Do Calculus

- Do-calculus: Contains, as subsets:
 - Backdoor criterion
 - Front-door criterion
- Allows analysis of more intricate structure beyond back- and front-door
- Uncovers **all** causal effects that can be identified from a given causal graph

Causal Inference: Packages and simulations

DoWhy tutorials:

<https://www.pywhy.org/dowhy/v0.14/>

<https://github.com/py-why/dowhy>

CausalGraphicalModels Tutorials:

<https://github.com/ijmbarr/causalgraphicalmodels>

Adjusting for the wrong variable: <http://www.degeneratestate.org/posts/2018/Jul/10/causal-inference-with-python-part-2-causal-graphical-models/>

Front-door: <http://www.degeneratestate.org/posts/2018/Sep/03/causal-inference-with-python-part-3-frontdoor-adjustment/>

Also see ML extensions to DoWhy, e.g. EconML:

<https://github.com/microsoft/EconML>