These are exercises for self-study and exam preparation. All material is examinable unless otherwise mentioned.

Exercise 1. *Factor analysis*

A friend proposes to improve the factor analysis model by working with correlated latent variables. The proposed model is

$$p(\mathbf{h}; \mathbf{C}) = \mathcal{N}(\mathbf{h}; \mathbf{0}, \mathbf{C}) \qquad p(\mathbf{v}|\mathbf{h}; \mathbf{F}, \boldsymbol{\Psi}, \mathbf{c}) = \mathcal{N}(\mathbf{v}; \mathbf{F}\mathbf{h} + \mathbf{c}, \boldsymbol{\Psi}) \tag{1}$$

where $\mathbf{C}$ is some covariance matrix, and the other variables are defined as in the lecture slides. $\mathcal{N}(\mathbf{x}; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes the pdf of a Gaussian with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$.

(a) What is marginal distribution of the visibles $p(\mathbf{v}; \boldsymbol{\theta})$ where $\boldsymbol{\theta}$ stands for the parameters $\mathbf{C}, \mathbf{F}, \mathbf{c}, \boldsymbol{\Psi}$?

(b) Assume that the singular value decomposition of $\mathbf{C}$ is given by

$$\mathbf{C} = \mathbf{E}\boldsymbol{\Lambda}\mathbf{E}^\top \tag{2}$$

where $\boldsymbol{\Lambda} = \text{diag}(\lambda_1, \ldots, \lambda_D)$ is a diagonal matrix containing the eigenvalues, and $\mathbf{E}$ is a orthonormal matrix containing the corresponding eigenvectors. The matrix square root of $\mathbf{C}$ is the matrix $\mathbf{M}$ such that

$$\mathbf{M}\mathbf{M} = \mathbf{C}, \tag{3}$$

and we denote it by $\mathbf{C}^{1/2}$. Show that the matrix square root of $\mathbf{C}$ equals

$$\mathbf{C}^{1/2} = \mathbf{E}\,\text{diag}(\sqrt{\lambda_1}, \ldots, \sqrt{\lambda_D})\mathbf{E}^\top. \tag{4}$$

(c) Show that the proposed factor analysis model is equivalent to the original factor analysis model

$$p(\mathbf{h}; \mathbf{I}) = \mathcal{N}(\mathbf{h}; \mathbf{0}, \mathbf{I}) \qquad p(\mathbf{v}|\mathbf{h}; \tilde{\mathbf{F}}, \boldsymbol{\Psi}, \mathbf{c}) = \mathcal{N}(\mathbf{v}; \tilde{\mathbf{F}}\mathbf{h} + \mathbf{c}, \boldsymbol{\Psi}) \tag{5}$$

with $\tilde{\mathbf{F}} = \mathbf{F}\mathbf{C}^{1/2}$, so that the extra parameters given by the covariance matrix $\mathbf{C}$ are actually redundant and nothing is gained with the richer parametrisation.

Exercise 2. *Independent component analysis*

(a) Whitening corresponds to linearly transforming a random variable $\mathbf{x}$ (or the corresponding data) so that the resulting random variable $\mathbf{z}$ has an identity covariance matrix, i.e.

$$\mathbf{z} = \mathbf{V}\mathbf{x} \quad \text{with} \quad \mathbb{V}[\mathbf{x}] = \mathbf{C} \quad \text{and} \quad \mathbb{V}[\mathbf{z}] = \mathbf{I}.$$

The matrix $\mathbf{V}$ is called the whitening matrix. We do not make a distributional assumption on $\mathbf{x}$, in particular $\mathbf{x}$ may or may not be Gaussian.

Given the eigenvalue decomposition $\mathbf{C} = \mathbf{E}\boldsymbol{\Lambda}\mathbf{E}^\top$, show that

$$\mathbf{V} = \text{diag}(\lambda_1^{-1/2}, \ldots, \lambda_d^{-1/2})\mathbf{E}^\top \tag{6}$$

is a whitening matrix.

©Gutmann, University of Edinburgh, 2026 CC BY 4.0

(b) Consider the ICA model

$$\mathbf{v} = \mathbf{A}\mathbf{h}, \qquad \mathbf{h} \sim p_{\mathbf{h}}(\mathbf{h}), \qquad p_{\mathbf{h}}(\mathbf{h}) = \prod_{i=1}^{D} p_h(h_i), \tag{7}$$

where the matrix $\mathbf{A}$ is invertible and the h_i are independent random variables of mean zero and variance one. Let $\mathbf{V}$ be a whitening matrix for $\mathbf{v}$. Show that $\mathbf{z} = \mathbf{V}\mathbf{v}$ follows the ICA model

$$\mathbf{z} = \tilde{\mathbf{A}}\mathbf{h}, \qquad \mathbf{h} \sim p_{\mathbf{h}}(\mathbf{h}), \qquad p_{\mathbf{h}}(\mathbf{h}) = \prod_{i=1}^{D} p_h(h_i), \tag{8}$$

where $\tilde{\mathbf{A}}$ is an orthonormal matrix.

©Gutmann, University of Edinburgh, 2026 CC BY 4.0